
Upper Mississippi River Basin


Nutrient Management Activities

June 2003

















Percentage of Wetland Losses in the Upper Mississippi River Basin States: 1780's to 1980's *





States

* Dahl. T.E. 1990 Wetlands Losses in the United States 1780's to 1980's.
U.S. Department of the Interior, Fish and Wildlife Service, Washington, D.C.,p6.




Grassland Losses

	Historic Acreage (ha)	Current Acreage (ha)	Decline
Illinois	8,900,000	900	99.9%
Indiana	2,800,000	404	99.9%
lowa	12,500,000	12,140	99.9%
Kansas	6,900,000	1,200,000	82.6%
Minnesota	7,300,000	30,350	99.6%
Missouri	5,700,000	30,350	99.5%



Root Systems of Prairie Plants





Upper Mississippi River Basin Current Landuses

 Agriculture 	67%
· Idle Agriculture	4%
• Urban	3%
 Forestry 	9%
 Federal 	12%
 Other 	5%



Longitudinal Changes in Mississippi River Floodplain Structure

Robert L. Delaney¹ and Mary R. Craig²




Figure 1. Mississippi River Segments

Table 1. Mississippi River Floodplain

River Segment	Approximate Floodplain Acres in 1,000s	Percent of Floodplain Behind Levees
Headwaters	328	<0.01%
Upper Mississippi (N)	496	3%
Upper Mississippi (S)	1,006	53%
Middle Mississippi	663	82%
Lower Mississippi	25,000	93%
Deltaic Plain	3,000	96%
TOTALS	30,493	90%



Habitat Alteration


River Segment	Floodplain Acreage (1000's)	% of Floodplain Behind Levees
Upper Mississippi (N)	496	3%
Upper Mississippi (S)	1,006	53%
Middle Mississippi	663	82%
Lower Mississippi	25,000	93%
Deltaic Plain	3,000	96%
Totals	30,493	90%



Ecological values of annual floodpulse



After a century of human alteration:





Post-dam: "chaos"

1930, Before Flooding

Terrestrial


Aquatic Vegetation

Open Water






1938



Aquatic Vegetation


Open Water





1965


- Terrestrial
- Aquatic Vegetation
- Open Water





1975


- Terrestrial
- Aquatic Vegetation
- Open Water






1993

- Terrestrial
- Aquatic Vegetation
- Open Water






OVERVIEW





Terrestrial
Aquatic Vegetation
Open Water

OVERVIEW





Terrestrial
Aquatic Vegetation
Open Water

Weaver Bottoms - 1950s





Weaver Bottoms - Current





US EPA'S Role


- Aiding in the identification of environmental problems
- Working in watersheds
- · Supporting the State's efforts




Aiding in the Identification of Environmental Problems

- Upper Mississippi River Data Base
- Fate and Transport Studies
- Further Watershed Analyses
- Pilot Projects
- Benefits of Existing Projects
















Pilot Projects

- Upper Halfway Creek (paired watershed)
- Wisconsin River Watershed (nutrient trading)
 - Baraboo River



Existing Projects

- Environmental Management Program (Water Resources Development Act 1986)
- · CREP, USDA
 - Minnesota River, MN
 - Illinois River, IL
 - Raccoon River, IA
- The Nature Conservancy Project
 - Spunky Bottoms
 - Emiquon
- Nutrient Farming
 - Hennipin Flood Control District

